

BLUEBERRY



	SOIL CONDITIONING	NUTRITION					
PRODUCT	REVITALIZE + FISH EMULSION	FULFIL	CALMATE	MICRONUTRIENTS	BEYOND FOLIAR 3-10-15	AGRODEX CAB	AGRODEX K35
APPLICATION RATES	2.5L/HA 5-10L/HA	1L/HA	40L/HA	2.5L/HA	10L/HA	5L/HA	8L/HA
GROWTH STAGE							
BUD BURST	○		APPLY WITH CALCIUM OR POTASSIUM NITRATE THROUGH IRRIGATION. MAXIMUM 40L/HA PER SEASON	○			
FIRST FLOWER		○		○	○	○	
2 WEEKS POST FLOWER		○			○		
4 WEEKS POST FLOWER					○	○	○
8 WEEKS POST FLOWER							○
2 WEEKS PRE HARVEST							○

- Shows application of product and timing
- Shows two applications of product during this growth period.

NOTES

- Early harvesting varieties can be coaxed into extended production as long as Agrodex K35 and Fulfill applications are kept up.
- Later varieties will come in slightly earlier.
- Tissue analysis should be sought to monitor nutrient status.
- Agrodex K35– Applications at bud burst is for frost protection only, apply if needed.
- Adjust Plant Food rates according to plant vigor, and results of analysis.
- Adjust Agrodex K35 - Rates according to crop load.

This program is designed to augment a soil fertility program and is offered as a guide only and dependant on growing conditions i.e. soil, climate, disease etc. Observe correct spray procedures as factors such as temperature, humidity, water rates can effect results. Always check compatibility before mixing products. Grower assumes all responsibility. Monitor nutrient status through tissue and soil analysis.